

Water Quality Enhancement Activity– WQL10 – Plant a cover crop that will scavenge residual nitrogen



Enhancement Description

Plant a cover crop that will scavenge nitrogen left in the soil after the harvest of a previous crop. Suitable cover crops include those with at least a “Very Good” rating for scavenging nitrogen as documented in *“Managing Cover Crops Profitably, 3rd Edition”* (Sarrantonio, 1998), Chart 2 Performance & Roles, pg 67. Examples include cereal rye, barley, forage radish and sorghum sudan.

Land Use Applicability

Cropland.

Benefits

Planting an annual cover crop to scavenge residual nutrients from cropland after the harvest of a previous crop effectively utilizes residual nutrient resources to supply following crops with nutrients required to efficiently produce food, forage, fiber, and cover while minimizing environmental degradation.

Criteria

Implementation of this enhancement requires:

1. The cover crop selected shall have the growth rate and rooting depth required to scavenge excess nitrogen from the root zone of the previous crop. Suitable cover crops include those with at least a “Very Good” rating for scavenging nitrogen as documented in *Managing Cover Crops Profitably, 3rd Edition, Chart 2 Performance & Roles, pg 67*. Examples include cereal rye, barley, forage radish and sorghum sudan.
2. Timing of planting and seeding rates for cover crops shall follow the recommendations in the respective NRCS Field Office Technical Guide (FOTG).
3. The producer must have a current soil test (no more than 3 years old).
4. Nitrogen application rates for the crop following the cover crop must be reduced by at least 15% from the “Land Grant University (LGU) recommendations to account for the recycling of N by the cover crop.
5. The producer shall not increase soil surface disturbance over existing benchmark conditions.



United States Department of Agriculture
Natural Resources Conservation Service

2010 Ranking Period 2

Documentation Requirements

Documentation for each Treatment area (field) and year of this enhancement describing these items:

1. A map showing where the activities are applied
2. Cover crop species planted
3. Cover crop planting date
4. Cover crop seeding rate (bu/ac)
5. Annual crop planted
6. Nitrogen application rates/amounts for the annual crop
7. Treatment acres



Water Quality Enhancement Activity – WQL10 – *Plant an Annual Cover crop that will Scavenge Residual Nitrogen*

Reference:

- **340 – Cover Crop**
- **590 – Nutrient Management**
- **Managing Cover Crops Profitably, 3rd Edition**
<http://www.mn.nrcs.usda.gov/technical/ecs/agron/Cover%20Crop/Managing%20Cover%20Crops%20Profitably.pdf>

The following annual cover crops have the growth rate and rooting depth required to scavenge excess nitrogen from the root zone of the previous crop:

Annual Ryegrass*

Barley*

Cereal Rye

Oats*

Radish*

Rapeseed (or Canola)*

Sorghum-Sudangrass

Wheat*

An estimate of the amount of N that is scavenged from each of these crops is given in the *Managing Cover Crops Profitably* book, in the write-up about each specific crop.

* If planted after Sept. 15, these crops will scavenge a relatively small amount of N.

The USDA is an equal opportunity provider and employer.



Species	Full Seeding Rate	Seeding Depth (inches)	Seeding Date	Comments
Annual Ryegrass	15 – 20 lbs/ac	¼ to ½	June 1-July 1 OR Aug 15 – Sept 15	Easily established. Good for use as overseeding row crop. May be seeded after harvest.
Barley	1.5 – 2 bu/ac	½ to 1 ½	Aug 15 - Sept 15	May be overseeded into growing crop or seeded after harvest.
Oats	1 – 2 bu/ac	½ to 1 ½	Aug 15 - Sept 15.	Can be seeded on rough plowed land (usually before Sept. 1) and will not need plowing the following spring
Cereal Rye	1 – 1 ¼ bu/ac	½ to 1 ½	Aug 15 - Sept 15	Easily established. Rapid growth in fall and spring. Has an allelopathic property.
Cereal Rye	¼ - ½ bu/ac	½ to 1 ½	Aug 15 - Sept 15	Use this rate only for cropland going into sugarbeets the following spring
Radish	8 – 10 lb/acre	¼ to ½	Aug 1 – Sept 15	Rapid fall growth, potential to capture nitrogen in large amts and from deep in the profile.
Rapeseed	5 lbs/acre	¼ to ¾	Aug 1- Sept. 15	Known for rapid fall growth, biomass production and nutrient scavenging.
Winter Wheat	1- 1 ½ bu/ac	½ to 1 ½	Aug 15 - Sept 15	Advantageous if site is seeded after Sept. 1 or under dry soil conditions.
Sorghum – Sudan	25 – 30 lbs/ac	½ to 1	May 15 - July 1	Advantageous to use on well drained and droughty sites.

Please Note:

1. Care is advised when determining the amount of N that will be available to the crop following the cover crop. Per acre yield in Minnesota may not be substantial depending on planting date and time of senescence or killing frost.
2. The lbs. per acre yield (dried) must be calculated; the percent N in plants just before death must be figured; and the percentage that can become available to the upcoming crop must be estimated (Yield X %N X % mineralized = available N).

The USDA is an equal opportunity provider and employer.



3. Assume that 50% of the N in the cover crop will be available to the upcoming crop in a plow down or incorporation system and 25% will be available in a system where the cover crop residues are left on the soil surface.
4. Nutrient application rates shall be consistent with University of Minnesota or contiguous land grant university recommendations.
<http://www.extension.umn.edu/CommodityCrops/>
<http://www.extension.umn.edu/distribution/cropsystems/DC5886.html>
5. Full Seeding Rates should be multiplied by the percent desired if mixtures are used.

The USDA is an equal opportunity provider and employer.