

Water Quality Enhancement Activity – WQL18 – Non-chemical pest control for livestock



Enhancement Description

Apply management techniques, devices, and biological agents that control external pests and internal parasites of livestock without the use of synthetic pesticides.

Land Use Applicability

Pastureland, rangeland and forestland (if the forest is grazed)

Benefits

Environmental benefits will be operation specific. Benefits may include, but are not limited to improved animal health, reduced risk to humans and improved water quality.

Pests and parasites can have a significant impact on the economic viability of livestock operations by affecting the performance and health of animals. The use of synthetic chemical treatments poses risks to water quality through animal contact and runoff. Farm workers are also exposed by handling chemicals and livestock. An alternative non-chemical pest control option can address these concerns and provide adequate pest/parasite control in many situations. Non-chemical control may also require increased monitoring and enhanced management applications which can affect a higher overall level of management efficiency.

Criteria

1. Prepare a written plan addressing basic management considerations, including:
 - a. Pests/parasites to be controlled, including correct species identification
 - b. Monitoring process (jug traps, baited cards, on-livestock counts, fecal egg counts, FAMANCHA, etc) to determine when control is needed and to monitor control effectiveness
 - c. Sanitation, cleaning feed/hay sites, and manure removal to reduce breeding sites
 - d. Rotational grazing and how it will be used to disrupt pest life cycles, minimum residual forage height to reduce parasite ingestion
2. Incorporate two or more of the following applications into the plan as appropriate:
 - a. Release of wasps that are parasitic to flies or the release of hister beetles
 - b. Traps for house and stable flies such as fly tape, paper, ribbons, etc.
 - c. Traps for biting flies or face flies
 - d. Walk through fly traps for horn flies
 - e. Fly vacuums
 - f. Bug zappers



Water Quality Enhancement Activity – WQL18 – *Non-chemical pest control for livestock*

Livestock external pests include deer, horn, house, horse and stable flies. Lice, ticks, scab mange and other external pests exist but require pesticide treatment for effective control.

Internal parasites of livestock include stomach worms (roundworms and nematodes), tapeworms, lungworms, liver flukes, and coccidian. The barber pole worm, a roundworm, is responsible for most losses in ruminants. Horsebots are parasites to the horse and related stock.

Pest Control Methods

Sanitation is the most important step in controlling flies and internal parasites.

- Eliminate breeding areas. Both stable and house flies breed in moist areas; straw and manure; moist, spilled feed and malt; manure drains or leaking water cups; and decaying vegetation around the premises. Stable flies are believed to overwinter in manure.
- Keep undisturbed manure as dry as possible. Along feed bunks, under fences, around waterers, and between pens or alleyways are ideal fly breeding areas.
- Keep waterers in good repair to avoid leaks. Grad lots and alleys to provide good drainage.
- Scrape empty pens, alleys, and fence lines to remove as much manure as possible and to improve drainage.
- Clean up spilled feed, decaying hay and silage.
- Control weeds to remove fly nesting areas.
- Certain manure management practices provide for manure storage under cages of poultry houses for long periods. If this is done in a manner where manure is subject to constant drying (which renders it nearly uninhabitable by flies), then the fly problem will be kept to a minimum.
- Where practical, spread manure thinly over fields under drying conditions, or heavy applications should be disked or shanked as liquid manure into the soil within 24 hours.
- Keep entryways clear of garbage cans or other fly-attracting material. Keep garbage or organic refuse cans tightly covered.
- Light traps, baited traps, and sticky tapes are physical controls for flies in barns.

The USDA is an equal opportunity provider and employer.

On pasture, to limit fly production, avoid large accumulations of residue and moisture at feeding sites. Proper cleanup and removal of residue may be necessary. Clean and dispose of waste prior to the beginning of the fly production season.

Other practical sanitation methods on pasture include:

- Continual physical movement of feeder locations between feedings.
- Rolling hay out in different locations throughout the pasture.
- Avoid using poor quality or rotted hay that will not be eaten.
- Avoid overfeeding regardless of feeding method, to prevent trampling and soiling of hay. Grinding hay may help prevent sorting and decrease waste.
- Feeding locations should have adequate drainage to keep moisture from accumulating around the feeder.

Releasing parasites can be considered after a good manure management system is in place. Commercially available parasites attack house and stable flies.

- Parasitic wasps should be periodically released before and during the fly season but before fly population levels build up.
- Select the right parasite. The two main groups of parasites are *Spalangia* and *Muscidifurax*. *Spalangia* appears to be the appropriate parasite for both stable and house flies. *Muscidifurax* appears to only control stable flies.
- Use quality products. Viability and purity of the parasite product are important. Find a reputable supplier that understands livestock fly problems and check to ensure shipments are viable when they arrive for release.
- Timing parasite release in the spring is critical to successful fly management. Parasite release should start when stable flies are noticed bothering livestock.
- Release the right number of parasites. Reliable suppliers should provide a formula or guidelines to calculate the number of wasps to use. The wasp number is based on the number of animals present, with 20 to 50 parasites per animal being released.

Pasture Management. Animals that graze are being exposed to internal parasites and the likelihood of re-infection is greater than those animals kept in confinement. In barns, animals should be fed using feeders rather than directly from the ground to limit contamination as a result of contact to bedding or manure. Pasture management strategies for internal parasite control include stocking density, use of multiple species, deferred grazing schedules and safe or clean pastures.

The USDA is an equal opportunity provider and employer.

- Grazing management must be the primary method for sustainable control of internal parasites. If pastures are not overstocked, there may be little difficulty with internal parasites. Forcing livestock to graze close to the ground tends to increase the occurrence of ingesting the infective larval stage of the parasites.
- Maintaining a grazing forage height of 4 inches or more will lower the probability of parasite infection. Livestock will avoid manure piles and the forage surrounding them which helps to avoid eating the larvae.
- Pasture recovery periods should be planned carefully. Longer rest periods are needed to avoid parasite contamination. At least 30 days of rest period should be planned; six months is more effective at ensuring larval death. One strategy is to combine a grazing period, rest period, and haying before subsequent grazing.
- Delay grazing of a pasture until parasite numbers have declined. Young winter or spring lambs should not graze infected pastures from March to May. Young stock, during their first year of grazing, are most susceptible to infection. Whenever possible, these animals should graze pasture that was not grazed by the same species in the previous year.
- The risk of infection can be lowered by delaying access to pasture until the dew has lifted or grass has dried after a rain. Parasites stay closer to the plant base in dry conditions and are less likely to be ingested.
- Plant anti-parasitic forages such as Birdsfoot Trefoil and Chicory.
- Sheep and goats share the same parasites and should not be co-grazed for parasite control.
- Cattle or horses do not share the same parasites as sheep or goats and can be cogenerated or alternately grazed on the same pastures.
- Incorporate an annual pasture into the grazing system and drag the stubble before planting to perennial pasture.
- Put goats in a browse area when hot humid conditions favor the rapid life cycle of internal parasites.
- Safe and/or Clean pastures should be part of the grazing acres. Safe pastures are ones that have been used for hay, silage or small grains. Safe pastures carry some parasite load but if managed properly provide a good way of controlling infection. Clean pasture can be a new seeding grazed for the first time or a pasture grazed the previous year by only a different livestock species that does not have common parasites, for example cattle grazed the year before sheep.
- Graze young stock away from their mothers on pastures with low parasite levels.

Monitor

- Fecal egg counts to determine the level of internal parasite pasture contamination.

The USDA is an equal opportunity provider and employer.

- FAMACHA examination of the lower eyelid of sheep and goats to determine degree of anemia caused by barber pole worms.
- Observe manure pats for the presence of dung beetles responsible for decomposition of the manure.
- Observe animal behavior. Animals may bunch up to avoid flies. They may kick feet or run about excessively to escape buzzing and striking flies. Watch for excessive licking or scraping that may indicate mange or mites.
- Inspect animals. Look for affected skin areas that have been scraped on animals, elevated skin areas along the spine on cattle, manure laden tags of wool on sheep, wounds in the mouths of horses and the vent area of poultry for dried blood.

References

Broce, A.B., et al., Managing Stable Fly Production at Pasture Feeding Sites MF-2662, Kansas State University, January 2005.

Kerr, S. 2007. Emerging concepts in small ruminant parasite control. Oregon State University. Corvallis OR.

<http://smallfarms.oregonstate.edu/sfn/spring08parasitecontrol>

Gamroth, Mike. 2009. Pests in the Organic Pasture. Michigan State University Extension and Oregon State University publication. Available online at

www.extension.org/article/18677

Greene, G., P. Sloderbeck, and J. Nechols. Biological Fly Control for Kansas Feedlots MF-2223, Kansas State University, April 1998.

Walker, E. and J. Stachecki. 1996. Livestock Pest Management Category 1D: A training manual for commercial applicators. Michigan State University Pesticide Education Program Bulletin E-2601.

<http://www.pested.msu.edu/Resources/bulletins/E2601.html>

Wells, A. 1999. Integrated Parasite Management for Livestock. National Center for Appropriate Technology. www.attra.ncat.org